

# Flora of the Darwin Region

## VOLUME 1

P.S. Short & I.D. Cowie (eds)



CABOMBACEAE

I.D. Cowie

**greening the  
Territory**

A Territory Government initiative

National Library of Australia  
Cataloguing-in-publication entry (PDF):  
Flora of the Darwin Region Volume 1  
ISBN: 9781921519949 (PDF)  
Series: Northern Territory Botanical Bulletin No. 37  
Notes: Includes bibliographical references and  
index.  
Subjects: Botany – Northern Territory – Darwin  
region – Plants – Identification  
Other Authors/Contributors: Short, P.S.,  
Cowie, I.D., Northern Territory Department  
of Natural Resources, Environment, the Arts  
and Sport.  
Dewey Number: 581.994295  
ISSN: 0314-1810  
Date of Publication: March 2011

---

Cover: *Helicteres procumbens* (Benth.) Cowie, *ined.*  
Cover Artist: R. Walter  
Editors: P.S. Short and I.D. Cowie  
Illustrations: M. Osterkamp  
Typesetting: D. Bisa and M. Voukalis  
Photographic Editor: B.M. Stuckey  
Authors contributing to this publication:  
D.E. Albrecht  
Northern Territory Herbarium, Department of  
Natural Resources, Environment, the Arts and  
Sport, P.O. Box 1120, Alice Springs, N.T. 0871,  
Australia  
R.M. Barker  
State Herbarium of South Australia, Plant  
Biodiversity Centre, P.O. Box 2732, Kent Town,  
S.A. 5071, Australia  
I.D. Cowie; B. Crase\*; D.J. Dixon\*; C.R. Dunlop\*;  
R.K. Harwood\*; R.A. Kerrigan\*; G.J. Leach\*;  
C. Mangion\*; P.S. Short; G.M. Wightman\*  
Northern Territory Herbarium, Department of  
Natural Resources, Environment the Arts and

Sport, P.O. Box 496, Palmerston, N.T. 0831,  
Australia  
L.A. Craven  
Australian National Herbarium, Centre for  
Plant, Biodiversity Research, G.P.O. Box 1600,  
Canberra, A.C.T. 2601, Australia  
L.L. Forman† (deceased)  
Royal Botanic Gardens, Kew, Richmond,  
Surrey, TW9 3AB, UK  
B. Jackes  
James Cook University Herbarium, School of  
Marine and Tropical Biology, James Cook  
University, Townsville, Qld 4811, Australia  
L. Jessup  
Queensland Herbarium Mt Coot-tha Road,  
Toowong, Qld 4066, Australia  
J. Palmer  
Australian National Herbarium, G.P.O. Box  
1600, Canberra, A.C.T. 2601, Australia

\*Former employee NT Herbarium

---

This book is copyright. Apart from any fair dealing  
for the purpose of private study, research, criticism  
or review, as permitted under the Copyright Act, no  
part of this publication may be produced by any  
process whatsoever without the written permission  
of the publisher.

© Northern Territory Government

Publisher:

Northern Territory Herbarium  
Department of Natural Resources,  
Environment, the Arts and Sport  
P.O. Box 496  
Palmerston, N.T. 0831, Australia

Suggested citation for this article:

Cowie, I.D. (2011). Cabombaceae. *In* Short, P.S.  
& Cowie, I.D. (eds), *Flora of the Darwin Region*.  
(Northern Territory Herbarium, Department of  
Natural Resources, Environment, the Arts and  
Sport). Vol. 1, pp. 1–2.  
[http://www.nt.gov.au/nreta/wildlife/plants\\_  
herbarium/index.html](http://www.nt.gov.au/nreta/wildlife/plants_herbarium/index.html)

# CABOMBACEAE

I.D. Cowie

Perennial aquatic *herbs*, rooted on substrate. *Stems* elongate, leafy. *Leaves* dimorphic, alternate, opposite or whorled, floating or submerged; floating leaves alternate, peltate, circular or rhombic to elliptic; submerged leaves, when present, opposite or whorled, deeply dissected, initially at least trichotomously then dichotomously. *Flowers* bisexual, axillary, solitary, hypogynous. *Perianth segments* 6–8; petals 3 or 4, white, yellow or dark maroon. *Stamens* 3–6 or 12–18; anthers tetrasporangiate and dithecal; staminodes absent. *Gynoecium* of 1–18 free carpels; ovules 1–3, anatropous; style short, stigmas terminal or decurrent. *Fruit* of coriaceous, indehiscent follicles.

A family of two genera and about eight species, found in tropical to warm temperate areas. One naturalised species is found in the N.T.

Sometimes recognised as a member of the Nymphaeaceae but differs in free carpels and has distinctive vessels. Yoo *et al.* (2005) discussed relationships and the phylogeny of the family.

Taxonomic reference: Yoo *et al.* (2005).

## CABOMBA Aubl.

Submerged attached perennials. *Submerged leaves* opposite or whorled, dissected, sessile or shortly petiolate. *Floating leaves* few or absent, blade rhombic to elliptic, entire, peltate. *Flowers* floating, solitary, axillary. *Perianth* 6-lobed, free, in 2 whorls. *Stamens* 3–6. *Carpels* often 2–4.

A genus of about seven species native to the Americas.

### \**C. caroliniana* A. Gray

*Herb*, stems to *c.* 1.5 m long, red. *Submerged leaves* on rachis 6–11 mm long, orbicular in outline, *c.* 50 mm diam, deeply dissected, initially divided palmately into *c.* 5 rays, each ray subsequently with 3–7 dichotomous divisions, ultimate segments terete; *floating leaves* 3–4 at apex of each stem, on petioles 7–26 mm long; blades narrowly rhombic, 17–30 mm long. *Tepals* oblong-obovate, 8–11 mm long; outer tepals white, green at base; inner tepals shortly clawed, white, bright yellow and cordate-auriculate at base. *Stamens* 6, *c.* 2.5 mm long; anthers *c.* 1.7 mm long. *Carpels* 2 or 3; usually 3-ovulate; stigmas capitate. *Fruiting peduncles* curved. *Flowering*. Mar.–May. *Fruiting*: not recorded.

Fig. 1 (Cowie 7459); Pl. 1 (unvouchered).

Native to America, naturalised in parts of Qld, N.S.W. and the N.T. Recorded from Marlow's Lagoon near Palmerston and Darwin River. The species is used as an aquarium plant and it is reported to be a serious weed in other parts of the world.

*Cabomba caroliniana*




Fig. 1



**REFERENCES**

Yoo, M.-J., Bell, C.D., Soltis, P.S. & Soltis, D.E. (2005). Divergence times and historical biogeography of Nymphaeales. *Systematic Botany* 30: 693–704.



Pl. 1 *Cabomba caroliniana* (Photo: C.G. Wilson)